

Precision QED $\otimes$ QCD Resummation Theory for LHC Physics: IR-Improved Scheme for Parton Distributions, Kernels, Reduced Cross Sections with Shower/ME Matching

B.F.L. Ward, S. Joseph, S. Majhi

Department of Physics, Baylor University, Waco, TX 76798-7316, USA

S.A. Yost

Department of Physics, The Citadel, Charleston, SC 29409, USA

We summarize the recent progress in a new approach to precision LHC physics based on the IR-improved DGLAP-CS theory as it relates to a new MC friendly exponentiated scheme for precision calculation of higher order corrections to LHC physics in which IR singularities from both QED and QCD are canceled to all orders in α and in α_s simultaneously in the presence of rigorous shower/ME matching. We present the first MC data comparing the implied new showers themselves with the standard ones using the HERWIG6.5 MC event generator as a test case at LHC energies.

The advent of the LHC marks the era of precision QCD, by which we mean predictions for QCD processes at the total theoretical precision tag of 1% or better. Here, we summarize the elements of our approach to such precision calculations [1, 2, 3] and its recent applications in Monte Carlo(MC) event generator studies, which are still preliminary.

At such a precision as we have as our goal, issues such as the role of QED [4, 5] are an integral part of the discussion and we deal with this by the simultaneous resummation of QED and QCD large infrared(IR) effects, $QED \otimes QCD$ resummation [1] in the presence of parton showers, to be realized on an event-by-event basis by MC methods. In proceeding with our analytical discussion, we first review this approach to resummation and its relationship to those in Refs. [6, 7]; this review is followed by a summary of the attendant new IR-improved DGLAP-CS theory. We conclude with a summary of illustrative applications and results, including the sample MC data.

To put the discussion in the proper perspective, we note that the authors in Ref. [8] have argued that the current state-of-the-art theoretical precision tag on single Z production at the LHC is $(4.1 \pm 0.3)\% = (1.51 \pm 0.75)\%(QCD) \oplus 3.79(PDF) \oplus 0.38 \pm 0.26(EW)\%$.¹

Turning now to our analysis, we note that, in Refs. [1, 2, 3], we have derived the following expression for the hard cross sections in the SM $SU_{2L} \times U_1 \times SU_3^c$ EW-QCD theory

$$d\hat{\sigma}_{\text{exp}} = e^{\text{SUM}_{\text{IR}}(\text{QCED})} \sum_{n,m=0}^{\infty} \frac{1}{n!m!} \int \prod_{j_1=1}^n \frac{d^3 k_{j_1}}{k_{j_1}} \prod_{j_2=1}^m \frac{d^3 k'_{j_2}}{k'_{j_2}} \int \frac{d^4 y}{(2\pi)^4} e^{iy \cdot (p_1 + q_1 - p_2 - q_2 - \sum k_{j_1} - \sum k'_{j_2}) + D_{\text{QCED}}} \tilde{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m) \frac{d^3 p_2}{p_2^0} \frac{d^3 q_2}{q_2^0}, \quad (1)$$

where the new YFS-style [10] residuals $\tilde{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m)$ have n hard gluons and m hard photons and we show the final state with two hard final partons with momenta p_2, q_2 specified for a generic 2f final state for definiteness. The infrared functions $\text{SUM}_{\text{IR}}(\text{QCED}), D_{\text{QCED}}$ are defined in Refs. [1, 2, 3]. This is the simultaneous resummation of QED and QCD large IR effects. Eq.(1) is exact.

Our approach to QCD resummation is fully consistent with that of Refs. [6, 7] as follows. First, Ref. [11] has shown that the latter two approaches are equivalent. We show in Refs. [2, 3] that our approach is consistent with that of

¹Recently, the analogous estimate for single W production has been given [9] as $\sim 5.7\%$.

Refs. [6] by exhibiting the transformation prescription from the resummation formula for the theory in Refs. [6] for the generic $2 \rightarrow n$ parton process as given in Ref. [12] to our theory as given for QCD by restricting (1) to its QCD component.

We show in Refs. [2, 3] that the result (1) allows us to improve in the IR regime² the kernels in DGLAP-CS [15, 16] theory as follows, using a standard notation:

$$\begin{aligned}
P_{qq}(z) &= C_F F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} \left[\frac{1+z^2}{1-z} (1-z)^{\gamma_q} - f_q(\gamma_q) \delta(1-z) \right], \\
P_{Gq}(z) &= C_F F_{YFS}(\gamma_q) e^{\frac{1}{2}\delta_q} \frac{1+(1-z)^2}{z} z^{\gamma_q}, \\
P_{GG}(z) &= 2C_G F_{YFS}(\gamma_G) e^{\frac{1}{2}\delta_G} \left\{ \frac{1-z}{z} z^{\gamma_G} + \frac{z}{1-z} (1-z)^{\gamma_G} \right. \\
&\quad \left. + \frac{1}{2} (z^{1+\gamma_G} (1-z) + z(1-z)^{1+\gamma_G}) - f_G(\gamma_G) \delta(1-z) \right\}, \\
P_{qG}(z) &= F_{YFS}(\gamma_G) e^{\frac{1}{2}\delta_G} \frac{1}{2} \{ z^2 (1-z)^{\gamma_G} + (1-z)^2 z^{\gamma_G} \}.
\end{aligned} \tag{2}$$

These results are being implemented by MC methods as we exhibit already below. We turn next to illustrative results and implications.

Firstly, we note that the connection to the higher order kernels in Refs. [17] is done [2]. Secondly, in the NS case, we find that the $n=2$ moment is modified by $\sim 5\%$ when evolved with (2) from 2GeV to 100GeV with $n_f = 5$ and $\Lambda_{QCD} \cong .2\text{GeV}$, for illustration. Thirdly, we have been able to use (1) to resolve violation [18, 19] of Bloch-Nordsieck cancellation in ISR at $\mathcal{O}(\alpha_s^2)$ for massive quarks [20]. Fourthly, the threshold resummation implied by (1) for single Z production at LHC shows a 0.3% QED effect and agrees with known exact results in QCD – see Refs. [1, 21, 22]. Fifthly, we have a new scheme [3] for precision LHC theory: in an obvious notation,

$$\sigma = \sum_{i,j} \int dx_1 dx_2 F_i(x_1) F_j(x_2) \hat{\sigma}(x_1 x_2 s) = \sum_{i,j} \int dx_1 dx_2 F'_i(x_1) F'_j(x_2) \hat{\sigma}'(x_1 x_2 s), \tag{3}$$

where the primed quantities are associated with (2) in the standard QCD factorization calculus. Sixthly, we have [1] an attendant shower/ME matching scheme, wherein, for example, in combining (1) with Herwig [23], Pythia [24], MC@NLO [25] or new shower MC's [26], we may use either p_T -matching or shower-subtracted residuals $\{\hat{\beta}_{n,m}(k_1, \dots, k_n; k'_1, \dots, k'_m)\}$ to create a paradigm without double counting that can be systematically improved order-by order in perturbation theory – see Refs. [1]. Finally, we show in Fig. 1 the effects of using the new kernels in a new version of Herwig6.5, Herwig6.5-YFS, on both the shower and pion energy fraction and p_T spectra for $2 \rightarrow 2$ hard sub-processes at LHC energies – as expected, the IR-improved spectra are softer. We look forward to the further exploration and development of the results presented herein.

Acknowledgments

One of us (B.F.L.W) acknowledges helpful discussions with Prof. Bryan Webber and Prof. M. Seymour. B.F.L. Ward also thanks Prof. L. Alvarez-Gaume and Prof. W. Hollik for the support and kind hospitality of the CERN TH Division and of the Werner-Heisenberg Institut, MPI, Munich, respectively, while this work was in progress.

Work partly supported by US DOE grant DE-FG02-05ER41399 and by NATO Grant PST.CLG.980342.

²This should be distinguished from the also important resummation in parton density evolution for the $z \rightarrow 0$ regime, where Regge asymptotics obtain – see for example Ref. [13, 14]. This improvement must also be taken into account for precision LHC predictions.




Figure 1: Distributions of the ISR parton energy fraction (a), P_T^2 (b), π^+ energy fraction (c), and P_T^2 (d) for the $2 \rightarrow 2$ hard subprocess in HERWIG-6.5, with (blue/green) and without (red) IR-improvement – preliminary.

References

- [1] C. Glosser, S. Jadach, B.F.L. Ward and S.A. Yost, *Mod. Phys. Lett.* **A 19**(2004) 2113; B.F.L. Ward, C. Glosser, S. Jadach and S.A. Yost, in *Proc. DPF 2004*, Int. J. Mod. Phys. **A20** (2005) 3735; in *Proc. ICHEP04, vol. 1*, eds. H. Chen et al., (World. Sci. Publ. Co., Singapore, 2005) p. 588; B.F.L. Ward and S. Yost, preprint BU-HEPP-05-05, in *Proc. HERA-LHC Workshop*, CERN-2005-014; in *Moscow 2006, ICHEP, vol. 1*, p. 505; *Acta Phys. Polon.* **B38** (2007) 2395; arXiv:0802.0724, in press at *Proc. RADCOR07*, 2008, and references therein.
- [2] B.F.L. Ward, *Adv. High Energy Phys.* **2008** (2008) 682312 ; DOI:10.1155/2008/682312.
- [3] B.F.L. Ward, *Ann. Phys.* **323** (2008) 2147.
- [4] S. Haywood, P.R. Hobson, W. Hollik and Z. Kunszt, in *Proc. 1999 CERN Workshop on Standard Model Physics (and more) at the LHC*, CERN-2000-004, eds. G. Altarelli and M.L. Mangano, (CERN, Geneva, 2000) p. 122 ; H. Spiesberger, *Phys. Rev.* **D52** (1995) 4936 ; W.J. Stirling, "Electroweak Effects in Parton Distribution Functions", talk presented at ESF Exploratory Workshop, *Electroweak Radiative Corrections to Hadronic Observables at TeV Energies* , Durham, Sept., 2003 ; M. Roth and S. Weinzierl, *Phys. Lett.* **B590** (2004) 190; J. Blumlein and H. Kawamura, *Nucl. Phys.* **B708** (2005) 467; *Acta Phys. Pol.* **B33** (2002) 3719; W. J. Stirling et al., in *Proc. ICHEP04*, eds. H. Chen et al. (World Sci. Publ., Singapore, 2005) p. 527; A.D. Martin et al., *Eur. Phys. J.* **C39** (2005) 155, and references therein.
- [5] A. Kulesza et al., in *Proc. RADCOR07*, in press, 2008; A. Denner et al., *ibid.*, in press, 2008; S. Pozzorini et al., *ibid.*, in press, 2008; A. Denner et al., *Nucl. Phys.* **B662** (2003) 299; G. Balossini et al., arXiv:0805.1129, and references therein.
- [6] G. Sterman, *Nucl. Phys.* **B 281**, 310 (1987); S. Catani and L. Trentadue, *Nucl. Phys.* **B 327**, 323 (1989); *ibid.* **353**, 183 (1991).
- [7] See for example C. W. Bauer, A.V. Manohar and M.B. Wise, *Phys. Rev. Lett.* **91** (2003) 122001; *Phys. Rev.* **D70** (2004) 034014
- [8] N.E. Adam et al., *J. High Energy Phys.* **0805** (2008) 062.

- [9] N.E. Adam et al., *J. High Energy Phys.* **0809** (2008) 133.
- [10] D. R. Yennie, S. C. Frautschi, and H. Suura, *Ann. Phys.* **13** (1961) 379; see also K. T. Mahanthappa, *Phys. Rev.* **126** (1962) 329.
- [11] C. Lee and G. Sterman, *ibid.* **D75** (2007) 014022.
- [12] M. Abyat *et al.*, *Phys. Rev.* **D74** (2006) 074004.
- [13] B.I. Ermolaev, M. Greco and S.I. Troyan, *Pos DIFF2006* (2006) 036, and references therein.
- [14] G. Altarelli, R.D. Ball and S. Forte, *PoS RADCOR2007* (2007) 028.
- [15] G. Altarelli and G. Parisi, *Nucl. Phys.* **B126** (1977) 298; Yu. L. Dokshitzer, *Sov. Phys. JETP* **46** (1977) 641; L. N. Lipatov, *Yad. Fiz.* **20** (1974) 181; V. Gribov and L. Lipatov, *Sov. J. Nucl. Phys.* **15** (1972) 675, 938; see also J.C. Collins and J. Qiu, *Phys. Rev. D* **39** (1989) 1398 for an alternative discussion of the lowest order DGLAP-CS theory.
- [16] C.G. Callan, Jr., *Phys. Rev. D* **2** (1970) 1541; K. Symanzik, *Commun. Math. Phys.* **18** (1970) 227, and in *Springer Tracts in Modern Physics*, **57**, ed. G. Hoehler (Springer, Berlin, 1971) p. 222; see also S. Weinberg, *Phys. Rev. D* **8** (1973) 3497; and references therein.
- [17] E.G. Floratos, D.A. Ross, C. T. Sachrajda, *Nucl. Phys.* **B129**(1977) 66; *ibid.* **B139**(1978) 545; *ibid.* **B152** (1979) 493, 1979; A. Gonzalez-Arroyo, C. Lopez and F.J. Yndurain, *Nucl. Phys.* **B153** (1979) 161; A. Gonzalez-Arroyo and C. Lopez, *Nucl. Phys.* **B166** (1980) 429; G. Curci, W. Furmanski and R. Petronzio, *Nucl. Phys.* **B175** (1980) 27; W. Furmanski and R. Petronzio, *Phys. Lett.* **B97** (1980) 437; E.G. Floratos, C. Kounnas and R. Lacaze, *Nucl. Phys.* **B192** (1981) 417; R. Hamberg and W. Van Neerven, *Nucl. Phys.* **B379** (1992) 143; S. Moch, J.A.M. Vermaseren and A. Vogt, *Nucl. Phys.* **B688** (2004) 101; *ibid.* **B691** (2004) 129.
- [18] C. Di'Lierto, S. Gendron, I.G. Halliday, and C.T. Sachradja, *Nucl. Phys.* **B183**(1981) 223; R. Doria, J. Frenkel and J.C. Taylor, *ibid.* **B168**(1980) 93, and references therein.
- [19] S. Catani, M. Ciafaloni and G. Marchesini, *Nucl. Phys.* **B264**(1986) 588; S. Catani, *Z. Phys.* **C37** (1988) 357.
- [20] B.F.L. Ward, BU-HEPP-07-04, arXiv.org:0707.2101; *Phys. Rev. D* **78** (2008) 056001.
- [21] U. Baur, S. Keller and W.K. Sakumoto, *Phys. Rev. D* **57** (1998) 199; U. Baur, S. Keller and D. Wackerroth, *ibid.* **59** (1998) 013002; U. Baur *et al.*, *ibid.* **65** (2002) 033007; S. Dittmaier and M. Kramer, *Phys. Rev. D* **65** (2002) 073007; Z. A. Zykunov, *Eur. Phys. J.* **C3** (2001) 9, and references therein.
- [22] R. Hamberg, W. L. van Neerven and T. Matsuura, *Nucl. Phys.* **B359** (1991) 343; W.L. van Neerven and E.B. Zijlstra, *Nucl. Phys.* **B382** (1992) 11; *ibid.* **B680** (2004) 513; C. Anastasiou *et al.*, *Phys. Rev. D* **69** (2004) 094008, and references therein.
- [23] C. Corcella *et al.*, hep-ph/0210213.
- [24] T. Sjostrand *et al.*, hep-ph/0308153.
- [25] S. Frixione and B. Webber, *J. High Energy Phys.* **0206** (2002) 029; S. Frixione, P. Nason and B. Webber, *ibid.* **0308** (2003) 007; and references therein.
- [26] S. Jadach and M. Skrzypek, in these *Proceedings*, 2008; *Comput. Phys. Commun.* **175** (2006) 511; P. Stevens *et al.*, *Acta Phys. Polon.* **B38** (2007) 2379, and references therein.